

Longitudinal polarization of hyperons in high $p_{\perp}$ jets in singly polarized pp collisions at high energies

Xu Qing-hua, Liu Chun-xiu* and Liang Zuo-tang

Department of Physics, Shandong University, Jinan, Shandong 250100, China

Abstract

We calculate the longitudinal polarizations of hyperons in high $p_{\perp}$ jets in pp collisions in which one of the protons is longitudinally polarized at RHIC energies using different models for the spin transfer in fragmentation process. The results show that the measurements of these polarizations can be used to study the spin transfer in high energy fragmentation processes in general and to test the different models in particular. Our results show especially that the magnitude of the polarization of Λ is rather small whereas that of Σ^+ is considerably larger in the large rapidity region. The differences between the results from different pictures for Σ^+ polarizations is also much larger. Hence, the measurement of Σ^+ polarization should be more effective to distinguish between different models especially the SU(6) or the DIS picture for spin transfer in fragmentation processes.

*Present address: Institute of High Energy Physics, Beijing 100039, China

1. Introduction

Spin effects in fragmentation processes play an important role in understanding the spin physics in strong interaction. With the developments in experiments, much attention has been attracted to the studies of the different aspects in this connection (see, e.g. Ref.[1-23] and references therein). Data are now available from different lepton-induced reaction experiments [1-5] and many of the theoretical studies [6-20] also concentrate on those reactions. It should be emphasized that the spin effects in the fragmentation processes can also be studied in polarized hadron-hadron collisions by measuring the properties of the hadrons in high transverse momentum $p_{\perp}$ jets created in these reactions [21-23]. Here, because of the validity of factorization theorem, the hadrons can be regarded as the pure results from the fragmentation of the scattered quark whose polarization can be determined using perturbation theory in quantum chromodynamics (QCD) and the quark distribution functions which determine the polarization before the hard scattering.

Compared with those in the lepton-induced reactions, the study of hadrons in high $p_{\perp}$ jets in polarized hadron-hadron collisions has the following properties. (1) Since the involved hard scatterings are strongly interacting processes rather than the electroweak processes involved in the lepton-induced reactions, the corresponding cross section should in general be larger. Furthermore, the luminosity of the incoming proton beams can in general be made higher than that for leptons in particular much higher than that for neutrino. Hence, the statistics can be improved in particular compared with the lepton-induced reactions with neutrino beams. (2) There are many different hard subprocesses which contribute to the production of the high $p_{\perp}$ jets. In particular, we have a large contribution from gluon interactions. This, on the one hand, makes the study more complicated than those in lepton-induced reactions, on the other, makes it more interesting since we can use it to study not only quark but also gluon polarization and fragmentation which are even poorly studied yet. Hence, it is also very important to know whether there exist kinematic regions where gluon fragmentation is important or where quark fragmentation dominates. In this case, we can

study them separately.

In this paper, we first make an analysis of the contributions of the different hard subprocesses to the production of different hyperons in different kinematic regions in pp collisions using a Monte Carlo event generator. Our results show that, there do exist kinematic regions where contributions from quark fragmentation dominate, while in the other gluon fragmentation plays an important role. We then calculate the longitudinal polarization of the hyperons in the regions where quark fragmentation dominates using the same method as we used in studying quark fragmentation in the lepton-induced reactions. In section 2, we present the results for the Monte Carlo analysis of the contributions of different hard processes. In section 3, we briefly summarize the calculations and the main results for the polarizations of the scattered partons from different hard subprocesses. In section 4, we present the results for the polarizations of different hyperons from high $p_\perp$ jets. A brief summary of the results is given in section 5.

2. Contributions of different hard subprocesses to hyperon production in high $p_\perp$ jets

We consider the inclusive production of hyperons with high transverse momentum $p_\perp$ in pp collisions at high energy. Here, the factorization theorem can be applied, hence the cross section can be written as²⁴,

$$E \frac{d\sigma}{d^3p}(pp \rightarrow HX) = \sum_{abcd} \int dx_a dx_b f_a(x_a, \mu^2) f_b(x_b, \mu^2) \frac{1}{\pi z_c} \frac{d\hat{\sigma}}{d\hat{t}}(ab \rightarrow cd) D_c^H(z_c, \mu^2). \quad (1)$$

Here, $f_a(x_a, \mu^2)$ and $f_b(x_b, \mu^2)$ are the unpolarized distribution functions of partons a and b in proton at the scale μ , x_a and x_b are the corresponding momentum fractions carried by a and b ; $D_c^H(z_c, \mu^2)$ is the fragmentation function of parton c into hadron H , z_c is the momentum fraction of parton c carried by the produced H ; $d\hat{\sigma}/d\hat{t}$ is the cross section at the parton level and $\hat{t}$ is the usual Mandelstam variable of the parton scattering. The cross section of the hard subprocess can be calculated using perturbative QCD. The summation in Eq. (1) runs over all possible subprocesses. To the leading order in pQCD, there are following subprocesses that contribute here, i.e. the scattering between quark(s) and/or antiquark(s),

$qq \rightarrow qq$, $q\bar{q} \rightarrow q\bar{q}$, $\bar{q}\bar{q} \rightarrow \bar{q}\bar{q}$; the scattering between gluons or that between gluon and quark, $qg \rightarrow qg$, $\bar{q}g \rightarrow \bar{q}g$, $gg \rightarrow gg$; the annihilation processes, $q\bar{q} \rightarrow gg$, $gg \rightarrow q\bar{q}$. The scattering matrix elements for these subprocesses can easily be calculated and the results can e.g. be found in Ref. [25]. For scattering angles less than $\pi/2$ in the center of mass frame of the scattering partons, the cross section for $gg \rightarrow gg$ is the largest, and $qg \rightarrow qg$ is the second largest one. They are much higher than the qq or $q\bar{q}$ scattering in particular at the scattering near $\pi/2$. Since the energies of gluons in protons are in general smaller compared with those for the quarks and/or antiquarks, we expect that gg scattering plays an important role for jets with moderately large transverse momenta $p_\perp$. But, for very high $p_\perp$, qq scattering should dominate.

Taking the fragmentation into account, we can estimate the contribution from each subprocess to the production of different hyperons. Presently, this can be conveniently done using Monte Carlo event generators which can give a good fit to the unpolarized data. We used PYTHIA^{26,27} in our calculations and obtained the results for Λ and Σ^+ as a function of pseudorapidity η in Figs.1 and 2. From these figures, we explicitly see that, at high $p_\perp$, e.g., $p_\perp > 8$ GeV at $\sqrt{s}=200$ GeV or $p_\perp > 13$ GeV at $\sqrt{s}=500$ GeV, only a few subprocesses dominate. The contributions of quark scatterings $qq \rightarrow qq$ (which denotes all the subprocesses in which the initial and final partons are all quarks) and $qg \rightarrow qg$ are important whereas that from $gg \rightarrow gg$ is suppressed significantly. We also calculated the contributions from quark jets or gluon jets. The results are shown in Figs.3 and 4. We see that the contribution from gluon fragmentation is negligibly small in this kinematic region, especially in the large rapidity region (e.g. $\eta > 1.5$). Therefore, the uncertainty of the calculations of hyperon polarizations in such kinematic regions caused by the unawareness of the polarized gluon fragmentation function should be very small. In our calculations which will be presented in Section 4, we will just neglect this contribution. However, at lower $p_\perp$, the contribution from $gg \rightarrow gg$ or that from gluon fragmentation is very important. As an example, we show in Fig. 5 the corresponding results for $p_\perp > 3$ GeV at $\sqrt{s}=500$ GeV. We see that the contribution of gluon fragmentation is higher than that of quark in the central

rapidity regions. Hence, we can study the spin transfer in quark fragmentation in the high $p_\perp$ region and that in gluon fragmentation in relatively small $p_\perp$ region.

3. Spin transfer and polarizations of the scattered quarks in different subprocesses

Since we are considering only the elementary subprocess with high $p_\perp$, the scattering matrix elements can be calculated to high precision using pQCD. The spin transfer factor for each subprocess can be obtained from the helicity amplitudes. The results show that, for all the different kinds of processes $ab \rightarrow cd$ (Here, a , b , c and d denote different partons, where a denotes the one from the longitudinally polarized incoming proton), the longitudinal polarization can be transferred to the outgoing parton c and d . The polarization transfer factors are in general different for different scattering processes. But they have a common feature that they are only functions of a single variable y , which is defined as $y \equiv k_b \cdot (k_a - k_c)/k_a \cdot k_b$, where k_a , k_b , k_c and k_d are the four-momenta of the partons a , b , c and d respectively. We summarize these results in Table 1.

From the table, we see the following: (i) Since we neglect quark mass, helicity conservation is valid. This can be seen from the results that $D_{(3)}^{a \rightarrow c} = D_{(5)}^{a \rightarrow c} = 1$, and that $D_{(4)}^{a \rightarrow c} = -D_{(4)}^{a \rightarrow d}$. (ii) In the scattering processes $qq' \rightarrow qq'$ and $qg \rightarrow qg$, the polarization transfer from one of the colliding objects to the other is the same, and they are also the same as that in the QED process $e^- \mu^- \rightarrow e^- \mu^-$ or $e^- q \rightarrow e^- q$. The result is a monotonously increasing function of y , which increases from zero to one as y goes zero to one. To get a feeling of all these spin transfer factors, we show them as functions of y in Fig. 6.

Multiplied by the polarizations of the incoming quarks or gluons, which are determined by the helicity distribution functions of the partons in proton, we obtain the polarizations of the outgoing quarks or gluons. They can be used to calculate the hyperon polarization produced in the fragmentation of these partons. Clearly, the obtained results depend on the polarized parton distribution functions we use. We will also show how strongly the final hyperon polarizations depend on the different sets of parameterization^{29–32} of these

distributions.

4. Longitudinal hyperon polarization in $pp \rightarrow HX$ in high $p_\perp$ regions

As we have seen in Sec. 2, for hyperons with considerably high transverse momenta, quark fragmentation dominates whereas gluon fragmentation can be neglected. In such kinematic regions, we can calculate the hyperon polarization using the same method as we used in treating $q_f^0 \rightarrow H_i + X$ in lepton-induced reactions. Measurements of the hyperon polarizations in these kinematic regions can be used as further tests of the different pictures for spin transfer in high energy quark fragmentation processes. The experimental studies can also be made in other kinematic regions where gluon fragmentation plays an important role. By comparing the obtained results with those obtained in the regions where quark fragmentation dominates, we can get useful information for gluon fragmentation. In this section, we present the calculations and the main results in the kinematic regions where quark fragmentation dominates.

A. The calculation method

The method of calculating the longitudinal polarization P_{H_i} of different hyperon H_i in the fragmentation of a longitudinally polarized quark q_f^0 was outlined in Ref. [15] using the inclusive process $e^+e^- \rightarrow H_i + X$ as an example. It has been applied to e^+e^- annihilations¹⁵ and deeply inelastic scattering using different lepton beams¹⁶. We now summarize the main points in the following not only for completeness but also for the following reasons: Here, we would like to emphasize what kinds of inputs that we need in the calculations and how we obtain these inputs. In this way, we can see from where the theoretical uncertainties originate in the calculations and how we should reduce these uncertainties by doing measurements for different hyperons and/or in different kinematic regions.

We consider $q_f^0 \rightarrow H_i + X$ and divide the produced H_i 's into the following groups:

(a) directly produced and contain the q_f^0 's; (b) decay products of heavier hyperons which were polarized before their decays; (c) directly produced but do not contain the q_f^0 's; (d) decay products of heavier hyperons which were unpolarized before their decays. Obviously, hyperons from (a) and (b) can be polarized while those from (c) and (d) are not. Hence, the polarization of H_i is given by,

$$P_{H_i} = \frac{\sum_f t_{H_i,f}^F P_f^{(q)} \langle n_{H_i,f}^a \rangle + \sum_j t_{H_i,H_j}^D P_{H_j} \langle n_{H_i,H_j}^b \rangle}{\langle n_{H_i}^a \rangle + \langle n_{H_i}^b \rangle + \langle n_{H_i}^c \rangle + \langle n_{H_i}^d \rangle}. \quad (2)$$

The different quantities here are defined and obtained in the following way:

(i) $P_f^{(q)}$ is the polarization of q_f^0 which is determined by the initial conditions and the elementary hard scattering processes.

(ii) $\langle n_{H_i,f}^a \rangle$ is the average number of H_i 's which are directly produced and contain q_f^0 of flavor f , and $\langle n_{H_i,H_j}^b \rangle$ is that from the decay of H_j 's which were polarized; P_{H_j} is the polarization of H_j ; $\langle n_{H_i}^a \rangle$, $\langle n_{H_i}^b \rangle$, $\langle n_{H_i}^c \rangle$ and $\langle n_{H_i}^d \rangle$ are the average numbers of H_i 's in group (a), (b), (c) and (d) respectively. These average numbers of the hyperons of different origins are determined by the hadronization mechanisms and should be independent of the polarization of the initial quarks. Hence, we can calculate them using a hadronization model which give a good description of the unpolarized data. We used Lund model²⁶ implemented by JETSET or LEPTO or PYTHIA²⁷ in our calculations for different reactions.

(iii) $t_{H_i,f}^F$ is the probability for the polarization of q_f^0 to be transferred to H_i in group (a), i.e. to the H_i which contains q_f^0 , and is called the polarization transfer factor, where the superscript F stands for fragmentation. It is taken as the fraction of spin carried by the f -flavor-quark divided by the average number of valence-quark of flavor f in H_i . This is the place where different pictures come in, since the contributions to the hyperon spin from different flavors are different in the SU(6) or the DIS picture^{11,15}. In the SU(6) picture, these contributions can be obtained from the SU(6) wave functions of the hyperons. In the DIS picture, for the $J^P = (1/2)^+$ octet hyperons, they are obtained from the DIS data on the spin dependent structure functions and those on hyperon decay. But, for the decuplet hyperons, they are unknown yet. In our calculations, we make a rough estimations

of these contributions by taking them as the same as those in the SU(6) picture. It will be interesting and important to study theoretically which picture is applicable in determining $t_{H_i,f}^F$. Discussions along this line can be found, for example, in [10] and the references given there.

(iv) t_{H_i,H_j}^D is the probability for the polarization of H_j to be transferred to H_i in the decay process $H_j \rightarrow H_i + X$ and is called decay polarization transfer factor, where the superscript D stands for decay. It is determined by the decay process and is independent of the process in which H_j is produced. For the octet hyperon decays, they are extracted from the materials in Review of Particle Properties²⁸. But, for the decuplet hyperons, we have to use an estimation based on the SU(6) quark model. (The details are given in e.g [7,15]). Hence, we see that decuplet hyperon decay is a major source of the theoretical uncertainties in our calculations of the final P_{H_i} 's in different reactions. For them, in the DIS picture, neither $t_{H_i,f}^F$ nor t_{H_i,H_j}^D is known yet. To reduce this uncertainty, it is important to consider the hyperons to which decay contributions are small.

We note also that $t_{H_i,f}^F$ and t_{H_i,H_j}^D should be considered as universal in the sense that they are the same in different reactions. The differences between different processes come from the polarization $P_f^{(q)}$ of the fragmenting quark q_f^0 and the average numbers $\langle n_{H_i,f}^a \rangle$, $\langle n_{H_i,H_j}^b \rangle$, $\langle n_{H_i}^c \rangle$, and $\langle n_{H_i}^d \rangle$ for hyperons H_i from the different origins. In pp collisions that we are considering in this paper, $P_f^{(q)}$ is determined by the polarization of the partons before the hard scattering and spin transfer factors in the scattering which are given and discussed in last section.

The average numbers $\langle n_{H_i,f}^a \rangle$, $\langle n_{H_i,H_j}^b \rangle$, $\langle n_{H_i}^c \rangle$, and $\langle n_{H_i}^d \rangle$ are determined by the hadronization mechanism and the relative abundance of different flavors that take part in the reactions. They should be independent of the process in which q_f^0 is produced and they are also independent of the polarization of q_f^0 . In pp collisions, as we can see from Eq.(1), these average numbers are determined by three factors, i.e., the parton distribution functions $f_a(x_a, \mu^2)$ and $f_b(x_b, \mu^2)$, the cross sections of hard subprocesses $d\hat{\sigma}/d\hat{t}$, and the fragmentation functions $D_c^H(z_c, \mu^2)$. The cross section $d\hat{\sigma}/d\hat{t}$ can be calculated with high accuracy from pQCD.

For $f_a(x_a, \mu^2)$, $f_b(x_b, \mu^2)$, and $D_c^H(z_c, \mu^2)$, although neither of them is theoretically clear yet, the form of the unpolarized parton densities in nucleon and that of the unpolarized fragmentation functions are empirically known to reasonably high accuracy. We can calculate them using the parameterizations of the structure functions and the phenomenological hadronization models. Since we are not concerned with the different correlations among the produced particles, different hadronization models lead essentially to the same results. Presently, these results can be obtained conveniently from the event generators using Monte Carlo method. We use the Lund string fragmentation model²⁶ implemented by PYTHIA²⁷ in our calculations for pp collisions.

B. Λ polarization in $pp \rightarrow \Lambda X$ with longitudinally polarized beam

Most of the studies now available on the spin transfer concentrate on Λ polarization since Λ is most abundantly produced among all the $J^P = (1/2)^+$ hyperons and its polarization can be measured easily through the decay channel $\Lambda \rightarrow p\pi^-$. Meanwhile, as we have seen already in e^+e^- annihilation¹⁵ and deeply inelastic lepton-nucleon scattering¹⁶, the origins of Λ are also the most complicated. They can be not only directly produced from the fragmentation of u , d , or s quark, but also from the decays of many different heavier hyperons, such as, Σ^0 , $\Xi^{0,-}$, $\Sigma^*(1385)$, and $\Xi^*(1530)$. In the large $p_\perp$ regions in pp collisions, the fragmentations of the quarks from the subprocesses $qg \rightarrow qg$ and $qq \rightarrow qq$ dominate. In particular for hyperons with large rapidities, the scatterings of quarks with large fractional momenta play the most important roles. Since at large x in proton, u quark density is much higher than that of d , and both of them are much higher than that of s , we expect that u and d , particularly u , quark fragmentation dominates the Λ production in the high $p_\perp$ and large rapidity region. However, u or d quark carries no spin of Λ in the SU(6) picture and only a small fraction of the spin of Λ in the DIS picture, we expect that, just as in deeply inelastic scattering¹⁶, the magnitudes of the P_Λ obtained in both models should be quite small, much smaller than those obtained in e^+e^- annihilation near the Z^0 pole. The influence of heavier hyperon

decay should be very important and the induced theoretical uncertainties in the calculations should be relatively large.

Using the event generator PYTHIA, we obtained the average numbers $\langle n_{\Lambda,f}^a \rangle$, $\langle n_{\Lambda,H_j}^b \rangle$, $\langle n_{\Lambda}^c \rangle$, and $\langle n_{\Lambda}^d \rangle$ for Λ production in $pp \rightarrow \Lambda X$. By inserting them into Eq. (2), we obtain P_{Λ} as a function of the pseudorapidity η . We show the different contributions to Λ production at $\sqrt{s} = 500$ GeV and $p_{\perp} > 13$ GeV as functions of η in Fig. 7 and P_{Λ} in Fig. 8. The moving direction of polarized beam proton is taken as the rapidity axis. From these results, we see in particular the following characteristics.

First, from Fig.7, we see that the contributions from u -quark fragmentation indeed dominate the Λ 's, in particular at large η . There are several direct consequences following from this : (i) the contributions from heavier hyperon decays are large, even larger than the directly produced contributions. To show this point more clearly, we present as an example in Fig. 9 the different contributions to P_{Λ} in the SU(6) picture. We see that the magnitudes of the contributions to P_{Λ} from Σ^0 and Σ^{*0} decays are both much larger than those from directly produced Λ 's. We see that even the qualitative tendency are influenced very much by these decay contributions. (ii) the resulting $|P_{\Lambda}|$ is relatively small, it is less than 3%. (iii) the differences between the results from SU(6) and those from the DIS picture are small.

Second, the contributions from those hyperons which are directly produced and contain the fragmenting q_f^0 's [origin (a) mentioned in section 4.A] or from the decays of the heavier hyperons which are directly produced and contain the fragmenting q_f^0 's [the type of heavier hyperons of origin (a)] are much larger than the corresponding contributions in the lepton-induced reactions. This is very interesting since these hyperons are all polarized and the polarizations are proportional to that of q_f^0 . In fact, in the limiting case that there are only these two types of hyperons contribute, we have,

$$P_{H_i}^{(lim)} = \frac{\sum_f t_{H_i,f}^F P_f^{(a)} \langle n_{H_i,f}^a \rangle + \sum_j t_{H_i,H_j}^D P_{H_j}^{(a)} \langle n_{H_i,H_j}^b \rangle}{\langle n_{H_i}^a \rangle + \langle n_{H_i}^b \rangle}. \quad (3)$$

Here, we use the superscript (a) in $P_{H_j}^{(a)}$ to denote the polarization of hyperon H_j of only origin (a). It is clear that, from Eq.(2), for such heavier hyperons,

$$P_{H_j}^{(a)} = \frac{\sum_f t_{H_j,f}^F P_f^{(q)} \langle n_{H_j,f}^a \rangle}{\langle n_{H_j}^a \rangle}. \quad (4)$$

By inserting it into Eq.(3), we obtain that,

$$P_{H_i}^{(lim)} = \frac{\sum_f [t_{H_i,f}^F \langle n_{H_i,f}^a \rangle + \sum_j t_{H_i,H_j}^D t_{H_j,f}^F \langle n_{H_j,f}^a \rangle R_{H_i,H_j}] P_f^{(q)}}{\langle n_{H_i}^a \rangle + \sum_j \langle n_{H_j}^a \rangle R_{H_i,H_j}}, \quad (5)$$

where R_{H_i,H_j} denotes the branching ratio for the decay $H_j \rightarrow H_i + X$. We see clearly that, in this limiting case, the P_{H_i} reflects more directly the polarization of the fragmenting quark q_f^0 . It is proportional to the $P_f^{(q)}$'s and the proportional constants depend on the picture for spin transfer and also the decay polarization transfer factors involved. We see clearly that P_Λ should depends rather strongly on $P_f^{(q)}$, which is determined by the spin transfer factors given in Table 1 and the polarized parton distribution functions. To see how sensitive it is, we used two different sets of parameterizations in the calculations. We see that the resulting difference in P_Λ is indeed quite significant.

C. Polarization of other hyperons

From the results that we obtained in the last subsection, we see that the contributions from the decay of heavier hyperons to Λ are large and the Λ polarization is very small. Since the contributions of different flavors to the spins of the $J^P = (3/2)^+$ hyperons are unknown yet in the DIS picture and only models are known to calculate the spin transfer in their decay processes, there are considerably large theoretical uncertainties in the results that we can obtain yet. On the other hand, as we have already seen^{15,16} in $e^+e^- \rightarrow Z^0 \rightarrow HX$ and in $\mu^-p \rightarrow \mu^- HX$, the origins of other $J^P = (1/2)^+$ hyperons, i.e., Σ and Ξ , are rather simple. The decay contributions are much smaller than the corresponding contributions for Λ production. The theoretical uncertainties in this case can be significantly reduced. In particular in the large rapidity region, the u quark fragmentation plays the dominant role and u carried most of the hyperon Σ^+ 's spin. Hence, the resulting Σ^+ polarization in $pp \rightarrow \Sigma^+ X$ should be much larger than that for Λ . Furthermore, also because of the

dominance of the u quark fragmentation, the production rate of Σ^+ should be comparable with that of Λ , which implies that the statistics for studying Σ^+ should be similar to that of Λ .

Using the event generator PYTHIA, we calculate the different contributions to the inclusive Σ^+ production in pp collisions. Using Eq.(2), we obtain P_{Σ^+} as a function of η at $p_{\perp} > 13$ GeV. The results are shown in Figs. 10 and 11 (a). Here, we see that the decay contribution from heavier hyperons is very small. It takes only a few percents for $p_{\perp} > 13$ GeV at $\sqrt{s}=500$ GeV. We see also that the obtained P_{Σ^+} is much larger than P_{Λ} and the difference between those results based on the two different pictures are much more significant, in particular for large η . From these calculations, we also see again that the contributions from the directly produced Σ^+ 's which contain the fragmenting quark play the dominate role. In fact, they give more than 75% of the whole produced Σ^+ at $p_{\perp} > 13$ GeV. In particular for $\eta > 1.5$, those directly produced and contain the fragmenting u -quarks take more than 93%. If we neglect all other contributions, we obtain the P_{Σ^+} in this limiting case, i.e. the P_{Σ^+} for those only from the origin (a) as,

$$P_{\Sigma^+}^{(lim)} = P_{\Sigma^+}^{(a)} = t_{\Sigma^+,u}^F P_u^{(q)}. \quad (6)$$

We see that it is directly proportional to the polarization of the u -quark, $P_u^{(q)}$, after the hard scattering and the proportional constant is just the fragmentation spin transfer $t_{\Sigma^+,u}^F$. This provides indeed an excellent place to test different pictures for spin transfer in fragmentation processes. It may provide also a nice tool to study the polarization of the quarks. Further studies along this line are under way³³. From Fig.11, we see also that the results for P_{Σ^+} from two different sets of parameterizations of the helicity parton distributions are rather small. This is because P_{Σ^+} depends mainly on Δu , which is essentially the same in the two sets of parameterizations.

We also make similar calculations for other hyperons, i.e., Σ^- , Ξ^0 and Ξ^- . The results are shown in Fig. 11 (b)-(d). We see that, $|P_{\Sigma^-}|$ exhibits similar behavior as $|P_{\Sigma^+}|$ but has opposite sign. This is because, similar to Σ^+ where u -quark fragmentation dominates, d -

quark fragmentation dominates Σ^- production. But Δd has opposite sign as Δu in proton. Both P_{Ξ^0} and P_{Ξ^-} are determined mainly by the fragmentation of s -quark but with some influence from u or d . Since u or d contributes negatively to the spin of Ξ , and Δu is positive in proton but Δd is negative, we should have a negative of u quark fragmentation to P_{Ξ^0} but a positive contribution from d to P_{Ξ^-} . This makes P_{Ξ^0} and P_{Ξ^-} behave a bit different. The difference can be seen in Fig.11. All these different features can be used to make further tests to the SU(6) or DIS picture used for the spin transfer in fragmentation processes.

5. Summary

In summary, after a Monte Carlo analysis of the different contributions, we calculated the longitudinal polarizations for different hyperons in high $p_\perp$ jets in pp collisions with longitudinally polarized beams using two different pictures: the SU(6) and the DIS pictures. The results show that the Λ polarization is rather small and has large contributions of different heavier hyperons decay. On the contrary, the contribution to Σ^+ production or that to Σ^- production is rather pure and the polarization is much larger. Hence, the measurements of these polarizations can give a good test to the validity of the two pictures. Such measurements can e.g. be carried out at RHIC in the near future.

Acknowledgments

We would like to thank Li Shi-yuan and Xie Qu-bing and other members in the theoretical particle physics group of Shandong University for helpful discussions. This work was supported in part by the National Science Foundation of China (NSFC) and the Education Ministry of China under Huo Ying Dong Foundation.

REFERENCES

1. ALEPH Collaboration, D. Buskulic *et al.*, Phys. Lett. B**374**, 319 (1996);
2. OPAL Collaboration, K. Ackerstaff *et al.*, Eur. Phys. J. C**2**, 49 (1998).
3. NOMAD Collaboration, P. Astier *et al.*, Nucl. Phys. B**588**, 3 (2000); *ibid* B605, 3 (2001).
4. HERMES Collaboration, A. Airapetian *et al.*, Phys. Rev. D**64**, 112005 (2001).
5. E665 Collaboration, M.R. Adams *et al.*, Euro. Phys. J. C**17**, 263 (2000).
6. R.L. Jaffe and Ji Xiang-dong, Phys. Rev. Lett. **67**, 552 (1991); Nucl. Phys. B**375**, 527 (1992).
7. G. Gustafson and J. Häkkinen, Phys. Lett. B**303**, 350 (1993).
8. K. Chen, G.R. Goldstein, R.L. Jaffe, and X. Ji, Nucl. Phys. B**445**, 380 (1995).
9. R.L. Jaffe, Phys. Rev. D**54**, R6581 (1996).
10. A.D. Adamov and G.R. Goldstein, Phys. Rev. D**56**, 7381 (1997); D**64**, 014021 (2001).
11. C. Boros and Liang Zuo-tang, Phys. Rev. D**57**, 4491 (1998).
12. A. Kotzinian, A. Bravar and D. von Harrach, Eur. Phys. J. C**2**, 329 (1998).
13. M. Anselmino, M. Boglione, U. D'Alesio, and F. Murgia, Eur. Phys. J. C**21**, 501(2001);
M. Anselmino, D. Boer, U. D'Alesio, and F. Murgia, Phys. Rev. D**63**, 054029 (2001).
14. B.Q. Ma, I. Schmidt and J.J. Yang, Phys. Rev. D**61**, 034017 (2000); B.Q. Ma, I. Schmidt, J. Soffer and J.J. Yang, Phys. Rev. D**62**, 114009 (2000); Phys. Rev. D**64**, 014017 (2001), erratum-*ibid*, D**64**, 099901 (2001).
15. Liu Chun-xiu and Liang Zuo-tang, Phys. Rev. D**62**, 094001 (2000).
16. Liu Chun-xiu, Xu Qing-hua and Liang Zuo-tang, Phys. Rev. D**64**, 073004 (2001).

17. Xu Qing-hua, Liu Chun-xiu and Liang Zuo-tang, Phys. Rev. **D63**, 111301(R) (2001).
18. Liang Zuo-tang, talk given at 31st Int. Sym. on Mitiparticle-dynamics (ISMD31), Datong, China, 1-9, Sep 2001, hep-ph/0111403.
19. Liang Zuo-tang, talk given at 3rd Circum-Pan-Pacific Symposium on High Energy Spin Physics, Beijing, China, 8-13, Oct 2001.
20. Xu Qing-hua *et al.*, talk given at 3rd Circum-Pan-Pacific Symposium on High Energy Spin Physics, Beijing, China, 8-13, Oct 2001.
21. D. de Florian, M. Stratmann, and W. Vogelsang, Phys. Rev. Lett. **81**, 530 (1998).
22. C. Boros, J. T. Londergan and A. W. Thomas, Phys. Rev. **D62**, 014021 (2000).
23. B.Q. Ma, I. Schmidt, J. Soffer and J. J. Yang, hep-ph/0107157.
24. J. Babcock, E. Monsay and D. Sivers, Phys. Rev. **D19**, 1483 (1979); N.S. Craigie *et al.*, Phys. Rep. **99**, 69 (1983)
25. R. K. Ellis, W. J. Stirling and B. R. Webber, “QCD and Collider Physics”, Cambridge University Press, (1996).
26. B. Andersson, G. Gustafson, G. Ingelman and T.Sjöstrand, Phys. Rep. **97**, 31 (1983).
27. T. Sjöstrand, Comp. Phys. Commun. **82**, 74 (1994).
28. Particle Data Group, D.E. Groom *et al.*, Euro. Phys. J. **C15**, 1 (2000).
29. M. Glück, E. Reya, M. Stratmann, and W. Vogelsang, Phys. Rev. **D63**, 094005 (2001).
30. M. Glück, E. Reya, and A. Vogt, Eur. Phys. J. **C5**, 461 (1998).
31. M. Glück, E. Reya, M. Stratmann, and W. Vogelsang, Phys. Rev. **D53**, 4775 (1996).
32. M. Glück, E. Reya, and A. Vogt, Z. Phys, **C67**, 433(1995).
33. Xu Qing-hua *et al.*, in preparation.

TABLES

Table 1. Polarization transfer factors $D_{(i)}^{a \rightarrow c}(y)$ and $D_{(i)}^{a \rightarrow d}(y)$ in the i -th kind of the elementary processes $ab \rightarrow cd$ when the incoming parton a is longitudinally polarized. Here, A and B are defined as $A \equiv 3 - 10y + 13y^2 - 6y^3 + 3y^4$ and $B \equiv 3 - 2y + y^2 - 2y^3 + 3y^4$.

The elementary process	The polarization transfer factors	
	$D_{(i)}^{a \rightarrow c}(y)$	$D_{(i)}^{a \rightarrow d}(y)$
$ab \rightarrow cd$		
(1) $q_f q_f \rightarrow q_f q_f$	$1 - 3y^4/A$	$(2y - 5y^2 + 6y^3)/A$
(2) $q_f \bar{q}_f \rightarrow q_f \bar{q}_f$	$1 - 3y^4/B$	$(2y - y^2 + 2y^3)/B$
(3) $q_f q_k(\text{or } \bar{q}_k) \rightarrow q_f q_k(\text{or } \bar{q}_k)$	1	$[1 - (1 - y)^2]/[1 + (1 - y)^2]$
(4) $q_f \bar{q}_f \rightarrow q_k \bar{q}_k$	$[(1 - y)^2 - y^2]/[(1 - y)^2 + y^2]$	$[y^2 - (1 - y)^2]/[(1 - y)^2 + y^2]$
(5) $q_f g \rightarrow q_f g$	1	$[1 - (1 - y)^2]/[1 + (1 - y)^2]$
(6) $g q_f \rightarrow g q_f$	1	$[1 - (1 - y)^2]/[1 + (1 - y)^2]$

FIGURES




Fig. 1. Contributions of the different parton level subprocesses to Λ production in $pp \rightarrow \Lambda X$.




Fig. 2. Contributions of the different parton level subprocesses to Σ^+ production in $pp \rightarrow \Sigma^+ X$.




Fig. 3. The contributions of quark and gluon jets to Λ production in $pp \rightarrow \Lambda X$.




Fig. 4. The contributions of quark and gluon jets to Σ^+ production in $pp \rightarrow \Sigma^+ X$.




Fig. 5. The contributions of quark and gluon jets to Λ production (a) and Σ^+ production (b) at $p_\perp > 3$ GeV in $pp \rightarrow HX$.




Fig. 6. Polarization transfer factors $D_{(i)}^{a \rightarrow c}(y)$ and $D_{(i)}^{a \rightarrow d}(y)$ in the i -th kind of the elementary processes $ab \rightarrow cd$ in Table 1.




Fig. 7. Different contributions to Λ production with $p_{\perp} > 13$ GeV in events originating from the d , u , or s quark fragmentation as a function of η in $pp \rightarrow \Lambda X$ at $\sqrt{s} = 500$ GeV. In (a)-(c), we see the five types of contributions from the fragmentation of d , u , or s quark respectively. The solid line denotes those which are directly produced and contain the fragmenting quark; the dashed, dash-dotted, *upper* dotted, *lower* dotted, denote those from the decay of Σ^0 , Ξ , Σ^* and Ξ^* which contain the fragmenting quark. In (d)-(f), we show the fractions R_i which are the ratios of the corresponding contributions to the sums of all different contributions.




Fig. 8. Longitudinal Λ polarization P_Λ as a function of η for Λ with $p_\perp > 13$ GeV in $pp \rightarrow \Lambda X$ at $\sqrt{s} = 500$ GeV with one of the beams is longitudinally polarized. ‘GRSV2000/GRV98’ denotes the Standard LO set of GRSV2000²⁹ for polarized distribution functions and GRV98³⁰ LO set for unpolarized distribution functions we used here; ‘GRSV96/GRV94’ denotes the Standard LO set of GRSV96³¹ and GRV94³² LO set correspondingly.




Fig. 9. Different contributions to P_Λ in the SU(6) picture for Λ with $p_\perp > 13$ GeV in $pp \rightarrow \Lambda X$ at $\sqrt{s} = 500$ GeV in the case that the Standard LO set of GRSV2000²⁹ for polarized distribution functions and GRV98³⁰ LO set for unpolarized distribution functions are used.




Fig. 10. Different contribution to Σ^+ production with $p_{\perp} > 13$ GeV in $pp \rightarrow \Sigma^+ X$ at $\sqrt{s} = 500$ GeV. The solid and dashed lines are respectively the contributions which are directly produced and contain the fragmenting quark u and s respectively; the dotted line corresponds to those which are originate from the decay of polarized heavier hyperons.




Fig. 11. Polarizations for Σ^+ , Σ^- , Ξ^0 and Ξ^- with $p_\perp > 13$ GeV in polarized $pp \rightarrow HX$ as a function of η at $\sqrt{s} = 500$ GeV. Two sets of polarized distribution functions as same as for Λ are used.

Table caption

Table 1: Polarization transfer factors $D_{(i)}^{a \rightarrow c}(y)$ and $D_{(i)}^{a \rightarrow d}(y)$ in the i -th kind of the elementary processes $ab \rightarrow cd$ when the incoming parton a is longitudinally polarized. Here, A and B are defined as $A \equiv 3 - 10y + 13y^2 - 6y^3 + 3y^4$ and $B \equiv 3 - 2y + y^2 - 2y^3 + 3y^4$.

Figure captions

Fig.1: Contributions of the different parton level subprocesses to Λ production in $pp \rightarrow \Lambda X$.

Fig.2: Contributions of the different parton level subprocesses to Σ^+ production in $pp \rightarrow \Sigma^+ X$.

Fig.3: The contributions of quark and gluon jets to Λ production in $pp \rightarrow \Lambda X$.

Fig.4: The contributions of quark and gluon jets to Σ^+ production in $pp \rightarrow \Sigma^+ X$.

Fig.5: The contributions of quark and gluon jets to Λ production (a) and Σ^+ production (b) at $p_\perp > 3$ GeV in $pp \rightarrow HX$.

Fig.6: Polarization transfer factors $D_{(i)}^{a \rightarrow c}(y)$ and $D_{(i)}^{a \rightarrow d}(y)$ in the i -th kind of the elementary processes $ab \rightarrow cd$ in Table 1.

Fig.7: Different contributions to Λ production with $p_\perp > 13$ GeV in events originating from the d , u , or s quark fragmentation as a function of η in $pp \rightarrow \Lambda X$ at $\sqrt{s} = 500$ GeV. In (a)-(c), we see the five types of contributions from the fragmentation of d , u , or s quark respectively. The solid line denotes those which are directly produced and contain the fragmenting quark; the dashed, dash-dotted, *upper* dotted, *lower* dotted, denote those from the decay of Σ^0 , Ξ , Σ^* and Ξ^* which contain the fragmenting quark. In (d)-(f), we show the fractions R_i which are the ratios of the corresponding contributions to the sums of all different contributions.

Fig.8: Longitudinal Λ polarization P_Λ as a function of η for Λ with $p_\perp > 13$ GeV in $pp \rightarrow \Lambda X$ at $\sqrt{s} = 500$ GeV with one of the beams is longitudinally polarized. ‘GRSV2000/GRV98’ denotes the Standard LO set of GRSV2000²⁹ for polarized distribution functions and GRV98³⁰ LO set for unpolarized distribution functions we used here; ‘GRSV96/GRV94’ denotes the Standard LO set of GRSV96³¹ and GRV94³² LO set correspondingly.

Fig.9: Different contributions to P_Λ in the SU(6) picture for Λ with $p_\perp > 13$ GeV in $pp \rightarrow \Lambda X$ at $\sqrt{s} = 500$ GeV in the case that the Standard LO set of GRSV2000²⁹ for polarized distribution functions and GRV98³⁰ LO set for unpolarized distribution functions are used.

Fig.10: Different contribution to Σ^+ production with $p_\perp > 13$ GeV in $pp \rightarrow \Sigma^+ X$ at $\sqrt{s} = 500$ GeV. The solid and dashed lines are respectively the contributions which are directly produced and contain the fragmenting quark u and s respectively; the dotted line corresponds to those which are originate from the decay of polarized heavier hyperons.

Fig.11: Polarizations for Σ^+ , Σ^- , Ξ^0 and Ξ^- with $p_\perp > 13$ GeV in polarized $pp \rightarrow HX$ as a function of η at $\sqrt{s} = 500$ GeV. Two sets of polarized distribution functions as same as for Λ are used.